

# Ultraperipheral collisions at the Large Hadron Collider

S. I. Godunov, E. K. Karkaryan, V. A. Novikov, A. N. Rozanov, M. I. Vysotsky,  
E. V. Zhemchugov

Lebedev Physical Institute  
Moscow, Russia

JINR, Dubna  
June 17, 2022


# Contents

- ▶ Ultraperipheral collisions [1806.07238].
- ▶ Survival factor [2106.14842].
- ▶ Long-lived charged particles [1906.08568].
- ▶ Semi-inclusive processes [to be published].

# Ultraperipheral collisions


[1806.07238]

# Ultraperipheral collisions at the LHC




Photon virtuality:  $Q^2 \equiv -q^2 = q_x^2 + q_y^2 + q_z^2 \ll (\gamma q_z)^2 \equiv \omega^2$


# Photon-photon processes studied at LHC



Nature Physics 13, 852 (2017)  
Phys.Lett. B797, 134826 (2019)




Phys.Lett. B777, 303 (2018)  
JHEP 1201, 052 (2012)



Phys.Rev. D94, 032011 (2016)  
JHEP 1608, 119 (2016)

# $p$ vs Pb




$$\sigma \sim Z^4$$

	$pp$	$Pb\ Pb$
Energy	13 TeV	5.02 TeV/(nucleon pair)
$Z$	1	82
$Z^4$	1	$4.5 \cdot 10^7$
Luminosity	$159\ fb^{-1}$ ratio: $6.6 \cdot 10^7$	$2.4\ nb^{-1}$
Duration	21 months (Run 2)	2 months (2015, 2018)
$\sqrt{s_{\gamma\gamma}}$	$\lesssim 2.6\ TeV$	$\lesssim 160\ GeV$


# $p$ vs Pb: forward detectors

[1909.10827]



Distance from the IP, m	200	420
$\xi$ range	0.015–0.15	0.002–0.02
6.5 TeV $p$ energy loss, GeV	97.5–975	13–130
in the center-of-mass frame, MeV	14–141	1.9–19
0.5 PeV $^{208}\text{Pb}$ energy loss, TeV	7.8–78	1.0–10
in the center-of-mass frame, GeV	2.9–29	0.37–3.7

# Equivalent photons approximation



$$\sigma(pp \rightarrow ppX) = \int_0^\infty d\omega_1 \int_0^\infty d\omega_2 \sigma(\gamma\gamma \rightarrow X) n_p(\omega_1) n_p(\omega_2)$$

$$n_p(\omega) = \frac{2Z^2\alpha}{\pi\omega} \int_0^\infty \left[ \frac{F_p(q_\perp^2 + (\omega/\gamma)^2)}{q_\perp^2 + (\omega/\gamma)^2} \right]^2 q_\perp^3 dq_\perp$$

$$Q^2 \equiv -q^2 = q_\perp^2 + \left( \frac{\omega}{\gamma} \right)^2$$

# Proton form factor

$$F_p(Q^2) = \frac{G_E(Q^2) + \tau G_M(Q^2)}{1 + \tau}$$

$$G_E(Q^2) = \frac{1}{\left(1 + \frac{Q^2}{\Lambda^2}\right)^2}$$

$$G_M(Q^2) = \frac{\mu_p}{\left(1 + \frac{Q^2}{\Lambda^2}\right)^2}$$

$$\tau = \frac{Q^2}{4m_p^2} \lesssim 0.01$$


$$\mu_p = 2.79$$

$$\Lambda_{\text{std.}}^2 = 0.71 \text{ GeV}^2$$

## Proton radius

$$r_p = 0.8414 \text{ fm} = \sqrt{12}/\Lambda$$

$$\Rightarrow \Lambda^2 = 0.66 \text{ GeV}^2.$$



[1307.6227]

# Proton EPA spectrum

Proton form factor with magnetic contribution taken into account:

$$F_p(Q^2) = \frac{G_E(Q^2) + \tau G_M(Q^2)}{1 + \tau},$$

$$G_E(Q^2) = \frac{1}{\left(1 + \frac{Q^2}{\Lambda^2}\right)^2}, \quad G_M(Q^2) = \frac{\mu_p}{\left(1 + \frac{Q^2}{\Lambda^2}\right)^2}$$

$$\tau = \frac{Q^2}{4m_p^2} \lesssim 0.01, \quad \Lambda^2 = 0.61 \text{ GeV}^2, \quad \mu_p = 2.79.$$

EPA spectrum:

$$\begin{aligned} n_p(\omega) = & \frac{\alpha}{\pi\omega} \left\{ \left(1 + 4u - 2(\mu_p - 1)\frac{u}{v}\right) \ln \left(1 + \frac{1}{u}\right) - \frac{24u^2 + 42u + 17}{6(u + 1)^2} \right. \\ & + \frac{\mu_p - 1}{(v - 1)^4} \left[ \frac{\mu_p - 1}{v - 1} (1 + 4u + 3v) - 2 \left(1 + \frac{u}{v}\right) \right] \ln \frac{u + v}{u + 1} \\ & + (\mu_p - 1) \frac{6u^2(v^2 - 3v + 3) + 3u(3v^2 - 9v + 10) + 2v^2 - 7v + 11}{3(u + 1)^2(v - 1)^3} \\ & \left. - (\mu_p - 1)^2 \frac{24u^2 + 6u(v + 7) - v^2 + 8v + 17}{6(u + 1)^2(v - 1)^4} \right\}, \\ u &= (\omega/\Lambda\gamma)^2, \quad v = (2m_p/\Lambda)^2. \end{aligned}$$

## $^{208}\text{Pb}$ form factor

Charge distribution is provided in form of Bessel spherical functions:

$$\rho(r) = \sum_{k=1}^N a_k j_0(k\pi r/R) \Theta(R - r); \quad j_0(x) = \sin(x)/x.$$

$a_k, R$  are numerical parameters.

Form factor is the Fourier transform of charge:

$$F_{\text{Pb}}(Q^2) = \frac{\int \rho(r) e^{i\vec{q}\vec{r}} d^3r}{\int \rho(r) d^3r} = \frac{\sin QR}{QR} \frac{\sum_{k=1}^N \frac{(-1)^k a_k}{(k\pi)^2 - (QR)^2}}{\sum_{k=1}^N \frac{(-1)^k a_k}{(k\pi)^2}}.$$


EPA spectrum:

$$n_{\text{Pb}}(\omega) = \frac{2Z^2\alpha}{\pi\omega} \int_0^\infty \left[ \frac{F(q_\perp^2 + (\omega/\gamma)^2)}{q_\perp^2 + (\omega/\gamma)^2} \right]^2 q_\perp^3 dq_\perp$$

Monopole approximation:

$$F_1(Q^2) = \frac{1}{1 + Q^2/\Lambda^2}, \quad n_1(\omega) = \frac{Z^2\alpha}{\pi\omega} \left[ (1 + 2u) \ln \left( 1 + \frac{1}{u} \right) - 2 \right], \quad u = \left( \frac{\omega}{\Lambda\gamma} \right)^2$$

# $^{208}\text{Pb}$ form factor




[1987]: At.Data and Nucl.Data Tabl. 36, 495 (1987)

[1995]: At.Data and Nucl.Data Tabl. 60, 177 (1995)

# $^{208}\text{Pb}$ charge density

$$\rho(r) = \sum_{k=1}^N a_k j_0(k\pi r/R) \Theta(R-r), \quad F(Q) = \frac{\int \rho(r) e^{i\vec{q}\cdot\vec{r}} d^3r}{\int \rho(r) d^3r}$$

	[1987]		[1995]	
	Quoted	Actual	Quoted	Actual
$Z = \int \rho(r) d^3r$	82	82.03	82	3806
$\langle r^2 \rangle^{1/2} = \left( \frac{\int \rho(r) r^2 d^3r}{\int \rho(r) d^3r} \right)^{1/2}, \text{ fm}$	5.499(1)	5.501	5.4785	8.0850




[1987]: At.Data and Nucl.Data Tabl. 36, 495 (1987)

[1995]: At.Data and Nucl.Data Tabl. 60, 177 (1995) ?


## EPA spectra

$E_p = 6.5 \text{ TeV}$ ,  $E_{\text{Pb}} = 522 \text{ TeV}$  (2.5 TeV/nucleon)



$$\hat{q}_p \gamma_p = 1.3 \text{ TeV}, \hat{q}_{\text{Pb}} \gamma_{\text{Pb}} = 79 \text{ GeV}$$

# CMS detector




Typical experimental requirements:  $|\eta| < \hat{\eta}$ ,  $p_T > \hat{p}_T$ .

$$\eta = -\ln \tan \frac{\theta}{2},$$

$$y = \frac{1}{2} \ln \frac{E_\chi + p_\chi \cos \theta}{E_\chi - p_\chi \cos \theta} \quad m_\chi \ll E_\chi \quad \eta$$

$$p_T = p_\chi \sin \theta$$



## Fiducial cross section

$$\frac{d\sigma_{\text{fid}}(pp \rightarrow pp\chi^+\chi^-)}{ds} = \int_{\max\left(\hat{p}_T, \frac{\sqrt{s}}{2 \cosh \hat{\eta}} \sqrt{1 - \frac{4m_\chi^2}{s}}\right)}^{\frac{\sqrt{s}}{2} \sqrt{1 - \frac{4m_\chi^2}{s}}} dp_T \frac{d\sigma(\gamma\gamma \rightarrow \chi^+\chi^-)}{dp_T} \frac{d\hat{L}}{ds}$$


Photon-photon luminosity:

$$\frac{d\hat{L}}{ds} = \frac{1}{4} \int_{-\hat{y}}^{\hat{y}} dy n\left(\frac{\sqrt{s}}{2} e^y\right) n\left(\frac{\sqrt{s}}{2} e^{-y}\right),$$

$\sqrt{s}$  — invariant mass of  $\chi^+\chi^-$  pair,  $y = \frac{1}{2} \ln \frac{\omega_1}{\omega_2}$  — its rapidity,

$$\hat{y} = \operatorname{asinh} \left[ \frac{\sqrt{s} p_T}{2(p_T^2 + m_\chi^2)} \left( \sinh \hat{\eta} - \sqrt{\cosh^2 \hat{\eta} + \frac{m_\chi^2}{p_T^2}} \cdot \sqrt{1 - \frac{4(p_T^2 + m_\chi^2)}{s}} \right) \right]$$


# ATLAS experiment: $pp \rightarrow pp\mu^+\mu^-$



Integrated cross section:

- ▶ Experiment:  $3.12 \pm 0.07$  (stat.)  $\pm 0.10$  (syst.) pb.
- ▶ Theory: 3.39 pb.

# ATLAS experiment: Pb Pb → Pb Pb $\mu^+ \mu^-$ (preliminary)



Integrated cross section:

$\sqrt{s}$ , GeV

- ▶ Experiment:  $32.3 \pm 0.3$  (stat.) $^{+4.0}_{-3.4}$  (syst.)  $\mu\text{b}$ .
- ▶ Theory:  $34.4 \mu\text{b}$  ( $\Lambda = 50 \text{ MeV}$ ) or  $57.4 \mu\text{b}$  ( $\Lambda = 80 \text{ MeV}$ ).

Survival factor


[2106.14842]

# Correction from strong interactions

Assuming only electromagnetic interactions:

$$n(\omega) = \frac{2Z^2\alpha}{\pi\omega} \int_0^\infty \left[ \frac{F(q_\perp^2 + (\omega/\gamma)^2)}{q_\perp^2 + (\omega/\gamma)^2} \right]^2 q_\perp^3 dq_\perp,$$

$$\sigma(pp \rightarrow ppX) = \int_0^\infty d\omega_1 \int_0^\infty d\omega_2 \sigma(\gamma\gamma \rightarrow X) n_p(\omega_1) n_p(\omega_2).$$




Including strong interactions:

$$n(\omega) = \int n(b, \omega) d^2b, \quad n(b, \omega) = \frac{Z^2\alpha}{\pi^2\omega} \left[ \int_0^\infty \frac{F(q_\perp^2 + (\omega/\gamma)^2)}{q_\perp^2 + (\omega/\gamma)^2} J_1(bq_\perp) q_\perp^2 dq_\perp \right]^2,$$

$$\begin{aligned} \sigma(pp \rightarrow ppX) \\ = \int_0^\infty d\omega_1 \int_0^\infty d\omega_2 \int d^2b_1 \int d^2b_2 \sigma(\gamma\gamma \rightarrow X) n_p(b_1, \omega_1) n_p(b_2, \omega_2) P(|\vec{b}_1 - \vec{b}_2|). \end{aligned}$$

$P(b)$  is the probability for the protons to survive after the collision with the impact parameter  $b$ .

# Polarization



$$\sigma(pp \rightarrow ppX) = \int_0^\infty ds \left[ \sigma_{\parallel}(\gamma\gamma \rightarrow X) \frac{dL_{\parallel}}{ds} + \sigma_{\perp}(\gamma\gamma \rightarrow X) \frac{dL_{\perp}}{ds} \right],$$

where

$$\frac{dL_{\parallel}}{ds} = \frac{1}{4} \int_{-\infty}^{\infty} dy \int d^2 b_1 \int d^2 b_2 n_p \left( b_1, \frac{\sqrt{s}}{2} e^y \right) n_p \left( b_2, \frac{\sqrt{s}}{2} e^{-y} \right) P(b) \cos^2 \varphi,$$

$$\frac{dL_{\perp}}{ds} = \frac{1}{4} \int_{-\infty}^{\infty} dy \int d^2 b_1 \int d^2 b_2 n_p \left( b_1, \frac{\sqrt{s}}{2} e^y \right) n_p \left( b_2, \frac{\sqrt{s}}{2} e^{-y} \right) P(b) \sin^2 \varphi$$

are photon-photon luminosities.

# Survival factor

$$\sigma(pp \rightarrow ppX)$$

$$= \int_0^\infty d\omega_1 \int_0^\infty d\omega_2 \int d^2 b_1 \int d^2 b_2 \sigma(\gamma\gamma \rightarrow X) n_p(b_1, \omega_1) n_p(b_2, \omega_2) P(|\vec{b}_1 - \vec{b}_2|)$$

vs


$$\sigma(pp \rightarrow ppX) = \int_0^\infty d\omega_1 \int_0^\infty d\omega_2 \sigma(\gamma\gamma \rightarrow X) n_p(\omega_1) n_p(\omega_2)$$

Neglecting the polarization,

$$S(s) = \frac{dL/ds}{dL/ds|_{P=1}},$$

where  $L = L_{\parallel} + L_{\perp}$ .

# Survival factor in $pp$ collisions




## Survival factor in Pb Pb collisions — no result

$$F_{\text{Pb}}(Q^2) = \frac{\sin QR}{QR} \frac{\sum_{k=1}^N \frac{(-1)^k a_k}{(k\pi)^2 - (QR)^2}}{\sum_{k=1}^N \frac{(-1)^k a_k}{(k\pi)^2}}$$

$$\begin{aligned} n(b, \omega) &= \frac{Z^2 \alpha}{\pi^2 \omega} \left[ \int_0^\infty \frac{F(q_\perp^2 + (\omega/\gamma)^2)}{q_\perp^2 + (\omega/\gamma)^2} J_1(bq_\perp) q_\perp^2 dq_\perp \right]^2 \\ &= \frac{Z^2 \alpha}{\pi^2 \omega R^4} \left[ \sum_{k=1}^K \frac{(-1)^k a_k}{(k\pi)^2} \right]^{-2} \times \\ &\quad \left[ \sum_{k=1}^K (-1)^k a_k \int_0^\infty \frac{x^2 \sin \sqrt{x^2 + y^2} J_1(cx)}{(x^2 + y^2 - (k\pi)^2)(x^2 + y^2)^{\frac{3}{2}}} dx \right]^2 \end{aligned}$$
$$x = q_\perp R, \quad y = R\omega/\gamma, \quad c = b/R.$$


# ATLAS experiment: $pp \rightarrow pp\mu^+\mu^-$



Integrated cross section:


- Experiment:  $3.12 \pm 0.07 \text{ (stat.)} \pm 0.10 \text{ (syst.) pb.}$
- No strong interactions: 3.39 pb.
- With strong interactions: 3.26 pb.

# ATLAS experiment: Pb Pb → Pb Pb $\mu^+\mu^-$ (preliminary)



# Long-lived charged particles

[1906.08568]



- ▶ Long-lived charged particles

- ▶ Live long enough to escape the detector (like muons).
- ▶ Usual search techniques:  $dE/dx$ , time of flight.
- ▶ Current bounds [1506.09173, 1609.08382, 1902.01636] are model-dependent.
- ▶ Example: SUSY chargino nearly degenerated with neutralino,  $m_\chi \gtrsim 100$  GeV.

- ▶ UPC approach:

- ▶ The particles leave tracks in the central detector allowing for reconstruction of their momenta  $\vec{p}_1, \vec{p}_2$ .
- ▶ Forward detectors provide the proton energies after the collision  $E_1, E_2$ .
- ▶ Collision kinematics is reconstructed. The mass of the particle


$$m = \sqrt{\frac{(2E_1E_2 + \vec{p}_1\vec{p}_2)^2 - \vec{p}_1^2\vec{p}_2^2}{4E_1E_2 + (\vec{p}_1 + \vec{p}_2)^2}}.$$

- ▶ Complementary to  $dE/dx$  or time of flight measurements.

- ▶ Background:


- ▶  $pp \rightarrow pp\mu^+\mu^-$  (and other processes producing muons).
- ▶ Pileup and diffractive scattering.

## Fermion pair production




Available integrated luminosity:  $pp - 159 \text{ fb}^{-1}$ ,  $Pb Pb - 2.4 \text{ nb}^{-1}$ .

## Fermion pair production (fiducial)




$p_T > 20$  GeV,  $|\eta| < 2.5$ , both protons hit the forward detectors.

# Expected signal (no pileup)




Assuming  $m_\chi = 100$  GeV and integrated luminosity 150/fb.

# Expected signal (with pileup)




Assuming pileup of 50 collisions at once.

# Expected signal (with pileup and longitudinal selection)




Longitudinal selection:  $|p_{||,1} + p_{||,2} - (E_1 - E_2)| < 20 \text{ GeV}$ .

# Expected signal (with pileup and longitudinal selection)



$$m_\chi = 190 \text{ GeV}.$$




Assuming integrated luminosity 150/fb.

# Semi-inclusive processes

[to be published]

## [2009.14537] measurement



Experimental selections:

- ▶  $p_T > 15 \text{ GeV}.$
- ▶  $|\eta| < 2.4.$
- ▶  $p_T^{\mu\mu} < 5 \text{ GeV}.$
- ▶  $20 \text{ GeV} < m_{\mu\mu} < 70 \text{ GeV}$  or  $m_{\mu\mu} > 105 \text{ GeV}.$
- ▶ At least one proton hits a forward detector.

$$\sigma(pp \rightarrow pX\mu^+\mu^-) = \sum_q \sigma(pq \rightarrow pq\mu^+\mu^-),$$

$$\text{For } \left(\frac{p_T^{\mu\mu}}{m_{\mu\mu}}\right)^2 \ll 1, \sigma(pq \rightarrow pq\mu^+\mu^-) = \int_0^\infty d\omega_1 \int_0^\infty d\omega_2 \sigma(\gamma\gamma \rightarrow \mu^+\mu^-) n_p(\omega_1) n_q(\omega_2)$$

$$n_q(\omega) = \frac{\alpha}{\pi\omega} \int_{\omega/E}^1 dx \int_{(\omega/x\gamma)^2}^{(p_T^{\mu\mu})^2} dQ^2 \frac{q_\perp^2}{Q^2} f_q(x, Q^2)$$

## Preliminary results

- ▶ Experiment:  $7.2 \pm 1.6$  (stat.)  $\pm 0.9$  (syst.)  $\pm 0.2$  (lumi.) fb.
- ▶ Exclusive process ( $pp \rightarrow pp\mu^+\mu^-$ ): 8.0 fb.
- ▶ Inclusive process ( $pp \rightarrow pX\mu^+\mu^-$ ): 8.7 fb.

Survival factor should reduce the calculated values by  $\sim 10\%$ . Errors are to be estimated yet.

# Conclusions

- ▶ Ultraperipheral collisions are a clean source of photon-photon collisions.
- ▶ Proton dipole form factor yields accurate results. Lead form factor is weird.
- ▶ Survival factor gives corrections of a few percents in the measurements considered.
- ▶ A new technique for detecting long-lived charged particles with the help of forward detectors. With  $150/\text{fb}$  of integrated luminosity, particles with lifetimes down to  $\sim 10^2$  ns and masses up to  $\sim 150$  GeV can be detected.
- ▶ Our calculation of semi-inclusive muon pair production cross section overshoots the measurement by a factor of 2.
- ▶ libepa (<https://github.com/jini-zh/libepa>) — A library for calculations of cross sections of ultraperipheral collisions under the equivalent photons approximation.

# Backup

# Proton EPA spectrum

$$F_p(Q^2) = \frac{G_E(Q^2) + \tau G_M(Q^2)}{1 + \tau},$$

$$G_E(Q^2) = \frac{1}{\left(1 + \frac{Q^2}{\Lambda^2}\right)^2}, \quad G_M(Q^2) = \frac{\mu_p}{\left(1 + \frac{Q^2}{\Lambda^2}\right)^2}, \quad \tau = -\frac{Q^2}{4m_p^2},$$

$$\begin{aligned} n_p(b, \omega) &= \frac{\alpha}{\pi^2 \omega} \left[ \int_0^\infty \frac{F_p(q_\perp^2 + (\omega/\gamma)^2)}{q_\perp^2 + (\omega/\gamma)^2} J_1(bq_\perp) q_\perp^2 dq_\perp \right]^2 \\ &= \frac{\alpha}{\pi^2 \omega} \left[ \frac{\omega}{\gamma} K_1\left(\frac{b\omega}{\gamma}\right) - \left(1 + \frac{(\mu_p - 1)\frac{\Lambda^4}{16m_p^4}}{\left(1 - \frac{\Lambda^2}{4m_p^2}\right)^2}\right) \sqrt{\Lambda^2 + \frac{\omega^2}{\gamma^2}} K_1\left(b\sqrt{\Lambda^2 + \frac{\omega^2}{\gamma^2}}\right) \right. \\ &\quad \left. + \frac{(\mu_p - 1)\frac{\Lambda^4}{16m_p^4}}{\left(1 - \frac{\Lambda^2}{4m_p^2}\right)^2} \sqrt{4m_p^2 + \frac{\omega^2}{\gamma^2}} K_1\left(b\sqrt{4m_p^2 + \frac{\omega^2}{\gamma^2}}\right) \right. \\ &\quad \left. - \frac{1 - \frac{\mu_p \Lambda^2}{4m_p^2}}{1 - \frac{\Lambda^2}{4m_p^2}} \cdot \frac{b\Lambda^2}{2} K_0\left(b\sqrt{\Lambda^2 + \frac{\omega^2}{\gamma^2}}\right) \right]^2, \end{aligned}$$

# $\gamma\gamma$ luminosities in $pp$ collisions

The probability for both protons to survive [hep-ph/0608271],

$$P(b) = \left(1 - e^{-\frac{b^2}{2B}}\right)^2, \quad B = 21 \text{ GeV}^2.$$

Photon-photon luminosities:

$$\begin{aligned} \frac{dL_{pp}^{\parallel}}{ds} &= \frac{\pi^2}{2} \int_{\hat{r}}^{\infty} b_1 db_1 \int_{\hat{r}}^{\infty} b_2 db_2 \int_{-\infty}^{\infty} dy n_p\left(b_1, \frac{\sqrt{s}}{2} e^y\right) n_p\left(b_2, \frac{\sqrt{s}}{2} e^{-y}\right) \\ &\times \left\{ 1 - 2e^{-\frac{b_1^2 + b_2^2}{2B}} \left[ I_0\left(\frac{b_1 b_2}{B}\right) + I_2\left(\frac{b_1 b_2}{B}\right) \right] + e^{-\frac{b_1^2 + b_2^2}{B}} \left[ I_0\left(\frac{2b_1 b_2}{B}\right) + I_2\left(\frac{2b_1 b_2}{B}\right) \right] \right\} \\ \frac{dL_{pp}^{\perp}}{ds} &= \frac{\pi^2}{2} \int_{\hat{r}}^{\infty} b_1 db_1 \int_{\hat{r}}^{\infty} b_2 db_2 \int_{-\infty}^{\infty} dy n_p\left(b_1, \frac{\sqrt{s}}{2} e^y\right) n_p\left(b_2, \frac{\sqrt{s}}{2} e^{-y}\right) \\ &\times \left\{ 1 - 2e^{-\frac{b_1^2 + b_2^2}{2B}} \left[ I_0\left(\frac{b_1 b_2}{B}\right) - I_2\left(\frac{b_1 b_2}{B}\right) \right] + e^{-\frac{b_1^2 + b_2^2}{B}} \left[ I_0\left(\frac{2b_1 b_2}{B}\right) - I_2\left(\frac{2b_1 b_2}{B}\right) \right] \right\} \end{aligned}$$

## Fiducial cross section

$$\frac{d\sigma_{\text{fid.}}(pp \rightarrow pp\chi^+\chi^-)}{ds} = \int_{\max\left(\hat{p}_T, \frac{\sqrt{s}}{2\cosh\hat{\eta}}\sqrt{1-\frac{4m_\chi^2}{s}}\right)}^{\frac{\sqrt{s}}{2}\sqrt{1-\frac{4m_\chi^2}{s}}} dp_T \left[ \frac{d\sigma_{||}(\gamma\gamma \rightarrow \chi^+\chi^-)}{dp_T} \frac{d\hat{L}^||}{ds} + \frac{d\sigma_{\perp}(\gamma\gamma \rightarrow \chi^+\chi^-)}{dp_T} \frac{d\hat{L}^{\perp}}{ds} \right]$$

Differential photon-photon cross sections:

$$\frac{d\sigma_{||}(\gamma\gamma \rightarrow \chi^+\chi^-)}{dp_T} = \frac{8\pi\alpha^2 p_T}{s(p_T^2 + m_\chi^2)} \cdot \frac{1 - \frac{2(p_T^4 + 2m_\chi^4)}{s(p_T^2 + m_\chi^2)}}{\sqrt{1 - \frac{4(p_T^2 + m_\chi^2)}{s}}}$$

$$\frac{d\sigma_{\perp}(\gamma\gamma \rightarrow \chi^+\chi^-)}{dp_T} = \frac{8\pi\alpha^2 p_T}{s(p_T^2 + m_\chi^2)} \cdot \frac{1 - \frac{2p_T^4}{s(p_T^2 + m_\chi^2)}}{\sqrt{1 - \frac{4(p_T^2 + m_\chi^2)}{s}}}$$

# Fiducial cross section

Fiducial luminosity:

$$\frac{d\hat{L}_{\parallel}}{ds} = \frac{1}{4} \iint d^2 b_1 d^2 b_2 P(b) \int_{-\hat{y}}^{\hat{y}} dy n\left(b_1, \frac{\sqrt{s}}{2} e^y\right) n\left(b_2, \frac{\sqrt{s}}{2} e^{-y}\right) \cos^2 \varphi$$

$$\frac{d\hat{L}_{\perp}}{ds} = \frac{1}{4} \iint d^2 b_1 d^2 b_2 P(b) \int_{-\hat{y}}^{\hat{y}} dy n\left(b_1, \frac{\sqrt{s}}{2} e^y\right) n\left(b_2, \frac{\sqrt{s}}{2} e^{-y}\right) \sin^2 \varphi$$

Rapidity cut:

$$\hat{y} = \operatorname{asinh} \left[ \frac{\sqrt{s} p_T}{2(p_T^2 + m_\chi^2)} \left( \sinh \hat{\eta} - \sqrt{\cosh^2 \hat{\eta} + \frac{m_\chi^2}{p_T^2}} \cdot \sqrt{1 - \frac{4(p_T^2 + m_\chi^2)}{s}} \right) \right]$$